

LIGHT PROPAGATION IN INHOMOGENEOUS UNIVERSE

Premana W. Premadi,¹ Hugo Martel,² and Richard A. Matzner^{3, 4}

1. Overview

Using a multi-plane lensing method that we have developed (Premadi 1996; Premadi, Martel, & Matzner 1998), we follow the evolution of light beams as they propagate through inhomogeneous universes. We use a P³M code to simulate the formation and evolution of large-scale structure. The resolution of the simulations is increased to sub-Megaparsec scales by using a Monte Carlo method to locate galaxies inside the computational volume according to the underlying particle distribution. The galaxies are approximated by isothermal spheres, with each morphological type having its own distribution of masses and core radii (Jaroszński 1992). The morphological types are chosen in order to reproduce the observed morphology-density relation (Dressler 1980). This algorithm has an effective resolution of 9 orders of magnitudes in length, from the size of superclusters down to the core radii of the smallest galaxies.

We consider cold dark matter models normalized to *COBE*, and perform a large parameter survey by varying the cosmological parameters Ω_0 , λ_0 , H_0 , and n (the tilt of the primordial power spectrum). The values of n are chosen by imposing particular values or σ_8 , the rms mass fluctuation at a scale of $8h^{-1}\text{Mpc}$. We use the power spectrum given in Bunn & White (1997). Table 1 gives the values of the parameters for all models. This is the largest parameter survey ever done in this field.

Table 1. Parameters for the various models

Ω_0	λ_0	H_0	n	σ_8	Ω_0	λ_0	H_0	n	σ_8
1.0	0.0	65.0	0.7234	0.9	0.2	0.0	65.0	1.3188	0.5
1.0	0.0	55.0	0.8465	1.0	0.2	0.0	75.0	1.2190	0.5
1.0	0.0	65.0	0.7698	1.0	0.2	0.0	75.0	1.2979	0.6
1.0	0.0	75.0	0.7094	1.0	0.2	0.0	75.0	1.3648	0.7
1.0	0.0	85.0	0.6605	1.0	0.7	0.3	65.0	0.7720	0.9
1.0	0.0	65.0	0.8120	1.1	0.7	0.3	75.0	0.7042	0.9
1.0	0.0	65.0	0.8506	1.2	0.7	0.3	65.0	0.8601	1.1
1.0	0.0	75.0	0.7893	1.2	0.7	0.3	75.0	0.7912	1.1
1.0	0.0	65.0	0.8861	1.3	0.5	0.5	65.0	0.7808	0.8
0.7	0.0	65.0	0.8461	0.9	0.5	0.5	75.0	0.7049	0.8
0.7	0.0	75.0	0.7773	0.9	0.5	0.5	65.0	0.8807	1.0
0.7	0.0	65.0	0.9346	1.1	0.5	0.5	75.0	0.8024	1.0
0.7	0.0	75.0	0.8648	1.1	0.2	0.8	65.0	0.9326	0.6
0.5	0.0	65.0	0.9457	0.8	0.2	0.8	75.0	0.8273	0.6
0.5	0.0	75.0	0.8686	0.8	0.2	0.8	65.0	1.0702	0.7
0.5	0.0	65.0	1.0439	1.0	0.2	0.8	55.0	1.2057	0.8
0.5	0.0	75.0	0.9656	1.0	0.2	0.8	65.0	1.0702	0.8
0.2	0.0	55.0	1.2187	0.3	0.2	0.8	75.0	0.9629	0.8
0.2	0.0	65.0	1.0966	0.3	0.2	0.8	85.0	0.8749	0.8
0.2	0.0	75.0	0.9993	0.3	0.2	0.8	65.0	1.1269	0.9
0.2	0.0	85.0	0.9191	0.3	0.2	0.8	65.0	1.1568	1.0
0.2	0.0	75.0	1.1228	0.4					

¹ Institute of Astronomy, Tohoku University Sendai 981, Japan

² Department of Astronomy, University of Texas Austin, TX 78712, USA

³ Center for Relativity, University of Texas Austin, TX 78712, USA

⁴ Department of Physics, University of Texas Austin, TX 78712, USA

2. The Ray Shooting Experiments

For each model, we perform numerous ray-tracing experiments, propagating beams of 341^2 light rays back in time up to redshifts $z = 3$ or 5 . Distance between two neighboring rays is about 1 arcsecond. This work is still in progress. So far, we have performed between 20 and 40 experiments for each model. Even with these small numbers of runs, we can already see significant differences between the various cosmological models. Figure 1 shows the individual contribution of each lens plane to the shear, as a function of the lens redshift, for sources located at $z = 3$ and $z = 5$, and for 3 cosmological models: An Einstein-de Sitter model, an open model with $\Omega_0 = 0.2$ and $\lambda_0 = 0$, and a flat model with $\Omega_0 = 0.2$ and $\lambda_0 = 0.8$ (these results are obtained by averaging over many experiments). The lenses planes contributing the most are located at intermediate redshifts, in spite of the fact that structures are more evolved at low redshifts, and galaxies are more crowded at high redshift. The shear peaks at lower redshift for the Einstein-de Sitter model than for the other two models. We interpret this effect as a consequence of the freeze-out of the growth of perturbations in models with $\Omega_0 < 1$. In such models, structures present at low redshift must have formed earlier than in the Einstein-de Sitter model. Hence, such models tend to have more structure at intermediate redshift for a given value of σ_8 .




Fig 1. Individual contribution of each lens plane to the shear vs redshift of the lens plane, for sources located at $z = 3$ (left panel) and $z = 5$ (right model). Solid curves: Einstein-de Sitter model; dotted curves: open model; dashed curves: flat, cosmological constant model.

Using these experiments, we can locate actual sources on the source plane, and compute their images on the image plane. In Figure 2, we plot the images of distant circular sources, to illustrate interesting cases that occurred in some experiments: magnification and shear (Fig. 2a), double image (Fig. 2b), Einstein ring (Fig. 2c), and triple image (Fig. 2d).

3. Preliminary Results

We can use the properties of the images to compute lensing statistics that can eventually be compared with observations. This is the ultimate goal of this project, and it will require many more experiments than the ones we have performed so far. For illustrative purposes, we computed, for various models, the probability that a lensed quasar will have multiple images. The results are presented in Table 2. As we see, the probabilities are small. Hence, most lensed quasars are simply magnified, with no image splitting. The results appear to vary among different cosmological models, but the number of runs is still too small to be statistically significant.




Fig. 2. Images of distant circular sources

Table 2. Multiple-image Probabilities

Ω_0	λ_0	H_0	σ_8	# or runs	Prob
1.0	0.0	65.0	1.0	37	0.040
0.2	0.0	65.0	0.5	38	0.088
0.5	0.0	65.0	1.0	30	0.054
0.5	0.0	75.0	1.0	20	0.070
0.7	0.0	65.0	1.1	18	0.059
0.5	0.5	65.0	1.0	20	0.089
0.2	0.8	65.0	0.8	39	0.130

References

Bunn, E. F., & White, M. 1997, *Ap.J.*, **480**, 6
Dressler, A. 1980, *Ap.J.*, **236**, 351
Jaroszński, M. 1992, *M.N.R.A.S.*, **255**, 655
Premadi, P. 1996, Ph.D. Thesis, University of Texas at Austin
Premadi, P., Martel, H., & Matzner, R. (1998), *Ap.J.*, **493**, 10